

PRACE GEOGRAFICZNE, zeszyt 107

Instytut Geografii UJ
Kraków 2000

Traute Crüger, Hans von Storch

DEVELOPMENT OF A DOWNSCALING MODEL FOR ESTIMATION OF AN 'ARTIFICIAL ICE CORE' DERIVED FROM LARGE SCALE PARAMETERS OF A 1000 YEAR GCM RUN

Abstract: Object of our work is the creation of 'artificial ice cores' by means of GCM data. Regression models for 3 Greenland ice core accumulation rates are presented which use the first 5 EOF time series of seasonal mean large scale 500 hPa stream function fields as predictors. The fitting data are the monthly NCEP Reanalysis data and real yearly ice accumulation time series from 1948 until 1992. The explained variance of the estimated ice accumulation lies between 56% and 74%. The regression model has been applied to a 1000 year coupled ocean/atmosphere control run (ECHAM4/HOPE) to estimate accumulation. The spectra of the real and the estimated accumulation are nearly 'white'. For the estimated accumulation there is a reduction of variance.

Key words: ice cores, reanalysis data, GCM data, linear regression.

1. Introduction

Many ice cores drilled from the Greenland and Antarctic ice sheet were taken to investigate features of the past climate. Much effort has also been done to perform GCM runs over hundreds of years, varying the underlying parameters like the CO₂ content or the solar radiation in order to eliminate the anthropogenic or natural impact on climate. Here, the ice accumulation rates are consistently estimated from GCM output and 'artificial ice cores' are constructed. These 'artificial ice cores' are compared to the real cores and used to estimate the natural variability and anthropogenic influence of ice accumulation rates. For the creation of 'artificial ice cores' one has to search for empirical relations between ice accumulation and GCM data. Since local features, especially precipitation is not well simulated by GCMs it is not possible to use this obvious predictor for our purpose (Fischer 1997). Therefore a statistical downscaling relation is derived which describes the local ice accumulation by means of well simulated

large scale atmospheric parameters such as geopotential, temperature or horizontal velocity fields. For developing a reliable downscaling relation the monthly NCEP Reanalysis data (Kalney et al. 1996) and three north and central Greenland ice core accumulation data sets from 1948 until 1992 are used (Fischer 1997; Schwager 1999). In Fig.1 the sites of the ice cores are shown. Their elevations lie between 2100 m (B21) and about 3000 m (B16, B29).

2. Regression Model: Statistical Technique, Parameters and Results

A simple downscaling method is a multiple linear regression using the EOF time series of the large scale field as predictors. This way the small scale noise is eliminated, while the main patterns are described by the EOF analysis, which are well simulated by GCMs. The EOF time series of the large scale field $G(x,t)$ result from a development in EOFs (v. Storch, Zwiers 1999):

$$G(x,t) = \sum_{i=1}^n G_i(x) A_i(t) \quad (1)$$

The $G_i(x)$ are the spatial patterns which are independent of time, whereas $A_i(t)$ are the EOF time series. Because the $A_i(t)$, $A_j(t)$ are orthogonal, the equation for multiple linear regression is:

$$A_{cc}(t) = K_1 A_1(t) + K_2 A_2(t) + \dots + K_n A_n(t) \quad (2)$$

With $A_{cc}(t)$ as local accumulation¹ and K_i as regression coefficients.

The three ice cores are described best using the first five EOF time series of the 500 hPa streamfunction. Using the temperature or the geopotential fields or combining them yields worse results in this set-up.

In the final form the three regression models use slightly different areas covering the northern hemisphere from the east American coast to east Asia in west/east direction and to the central Sahara in the south. The main differences of the models are the different periods which have to be considered by the regression models. B21 is best described using only the mean May/June field. The B16 model uses the mean streamfunction field from August until October and the B29 model from June until February.

Fig. 2 shows the ice accumulation anomalies for the real ice core and the estimated accumulation obtained by applying cross-validation (v. Storch, Zwiers 1999). The explained variance for the accumulation of B16 is 56%, of B21 69% and of B29 74%.

¹ Because of uncertainties of ice accumulation measurements and dating a weighted running mean (with weights 1-2-3-2-1) has been applied to the accumulation time series. In order to relate the yearly ice accumulation rates to the atmospheric data the date of the spring signal is set to May 1st.